

RESEARCH ARTICLE

Sensory Processing

## Reproductive state modulates utricular auditory sensitivity in a vocal fish

 Lorianze S. Rogers,<sup>1</sup> Allison B. Coffin,<sup>2</sup> and  Joseph A. Sisneros<sup>1,3,4</sup>

<sup>1</sup>Department of Psychology, University of Washington, Seattle, Washington; <sup>2</sup>Department of Integrative Physiology and Neuroscience, Washington State University, Vancouver, Washington; <sup>3</sup>Department of Biology, University of Washington, Seattle, Washington; and <sup>4</sup>Virginia Merrill Bloedel Hearing Research Center, University of Washington, Seattle, Washington

### Abstract

The plainfin midshipman, *Porichthys notatus*, is a seasonally breeding vocal fish that relies on acoustic communication to mediate nocturnal reproductive behaviors. Reproductive females use their auditory senses to detect and localize “singing” males that produce multiharmonic advertisement (mate) calls during the breeding season. Previous work showed that the midshipman sacculus, which is considered the primary end organ used for hearing in midshipman and most other fishes, exhibits reproductive state and hormone-dependent changes that enhance saccular auditory sensitivity. In contrast, the utricle was previously posited to serve primarily a vestibular function, but recent evidence in midshipman and related toadfish suggests that it may also serve an auditory function and aid in the detection of behaviorally relevant acoustic stimuli. Here, we characterized the auditory-evoked potentials recorded from utricular hair cells in reproductive and nonreproductive female midshipman in response to underwater sound to test the hypothesis that variation in reproductive state affects utricular auditory sensitivity. We show that utricular hair cells in reproductive females exhibit up to a sixfold increase in the utricular potential magnitude and have thresholds based on measures of particle acceleration (re:  $1 \text{ ms}^{-2}$ ) that are 7–10 dB lower than nonreproductive females across a broad range of frequencies, which include the dominant harmonics of male advertisement calls. This enhanced auditory sensitivity of the utricle likely plays an essential role in facilitating midshipman social and reproductive acoustic communication.

**NEW & NOTEWORTHY** In many animals, vocal-acoustic communication is fundamental for facilitating social behaviors. For the vocal plainfin midshipman fish, the detection and localization of social acoustic signals are critical to the species’ reproductive success. Here, we show that the utricle, an inner ear end organ often thought to primarily serve a vestibular function, serves an auditory function that is seasonally plastic and modulated by the animal’s reproductive state effectively enhancing auditory sensitivity to courting male advertisement calls.

auditory; hair cells; seasonal plasticity; utricle

### INTRODUCTION

Seasonal changes in sensory processing related to an animal’s reproductive cycle occur in many nonmammalian vertebrates including songbirds, amphibians, and fishes (for review see Refs. 1–3). Furthermore, reproductive-related changes in sensory processing of auditory information occur in a number of seasonally breeding species that rely on acoustic communication to mediate social interactions in a reproductive context [e.g., birds: (4–9); amphibians: (10–12); and fishes: (13, 14)]. However, previous work has primarily focused on reproductive state-dependent changes in sensitivity of the central auditory system (5, 6, 10–12) or primary

hearing organs of the peripheral auditory system (4, 13, 14). Here, we consider reproductive state-dependent changes in the frequency sensitivity and auditory gain of the utricle, an end organ not often associated with an auditory function, in a seasonally breeding vertebrate for which the detection and localization of conspecific acoustic signals are critical to its reproductive success.

The plainfin midshipman fish (*Porichthys notatus*) is a seasonally breeding vocal fish that produces social acoustic signals for intraspecific communication during the reproductive season. The social behaviors of this nocturnally active species are highly dependent upon the production and reception of acoustic signals, which makes the midshipman an excellent



model for investigating the neural mechanisms of acoustic communication, especially those related to seasonal changes in vocal-acoustic behavior and auditory reception (15–17). During the late spring and summer, midshipman migrate into the shallow intertidal zone to reproduce and care for their offspring. Courting (type I) males establish nest sites in the rocky substrate where they produce long-duration multiharmonic advertisement calls to attract gravid females for reproduction (18). Previous work has shown that females exhibit reproductive state- and hormone-dependent changes in the auditory sensitivity of the saccule, such that reproductive females are better suited than nonreproductive females to detect conspecific vocalizations (13, 19, 20). This steroid-, reproductive state-dependent modulation of auditory saccular sensitivity is thought to enhance the coupling of sender and receiver in the midshipman acoustic communication system.

In most fishes, the inner ear saccule is often the largest otolithic end organ and most associated with hearing (21, 22), whereas the smaller utricle has been posited to serve primarily a vestibular function as a gravistatic organ (23–27). However, recent evidence in toadfish and midshipman (Family Batrachoididae) suggests that the utricle is capable of detecting and encoding behaviorally relevant acoustic stimuli including conspecific vocalizations (28–30). Yet, the extent to which the utricle may exhibit reproductive-related changes in auditory sensitivity to social acoustic signals remains unknown.

Here, we test the hypothesis that seasonal variation in reproductive state modulates the auditory sensitivity of the utricle in female plainfin midshipman. We compare the auditory-evoked utricular potentials of reproductive and nonreproductive females to determine whether there are differences related to reproductive state in the frequency response and auditory threshold of utricular hair cells to behaviorally relevant auditory stimuli. We show that the utricle serves an auditory function that is seasonally plastic and highly adapted in reproductive females to detect the dominant frequencies of conspecific vocalizations.

## MATERIALS AND METHODS

### Animal Collection and Husbandry

Nonreproductive adult female plainfin midshipman fish, *Porichthys notatus* Girard 1854, were collected via otter trawls (R/V Kittiwake, Friday Harbor Laboratories) in January 2021 from the Puget Sound near Edmonds, WA, at depths ranging from 85 to 100 m. Reproductive adult female plainfin midshipman were collected during their breeding season (May–June 2021) by hand at low tide from exposed nest sites in the rocky intertidal area at Seal Rock near Brinnon, WA. Following collection, fish were transported to the University of Washington and housed in a 350 L recirculating artificial saltwater tank maintained at  $15 \pm 2^\circ\text{C}$  and kept on either a winter (9/15-h) or summer (12/12-h) light/dark photoperiod, which corresponds with the nonreproductive and reproductive ambient photoperiods, respectively. Before each physiology experiment, standard length (SL; cm) and body mass (BM; g) were recorded and sex was determined by visual inspection of the gonads. The gonadosomatic index {GSI;


defined here as  $100 \times [\text{gonad mass}/(\text{BM} - \text{gonad mass})]$  for each fish was recorded following each experiment. Utricular hair cell potential recordings were performed within 17 days after trawl collection in the winter and 14 days after hand collection during the summer to minimize any effects of prolonged captivity on midshipman auditory sensitivity while still allowing the animals to recover from capture-related stress.

### Acoustic Stimulus and Calibration

The methodology used for acoustic stimulus presentation and calibration was similar to that of previously published work (13, 29, 31–35). Acoustic stimuli were generated by a lock-in amplifier (SR830, Stanford Research Systems, Sunnyvale, CA), which sent pure tone signals to an audio amplifier (BG-1120, TOA Corporation, Hyogo, Japan) and then to an underwater speaker (UW-30, Telex Communications, Burnsville, MN). The midshipman utricle is likely highly sensitive to particle motion along the horizontal plane as both the otolith (i.e., lapillus) and hair cells are oriented in the horizontal plane (Fig. 1B, Supplemental Fig. S1; all Supplemental material is available at <https://doi.org/10.6084/m9.figshare.20363625>; but also see Fig. 6 in Ref. 36). Therefore, the underwater speaker was positioned such that the speaker's face resided along the horizontal plane and was fully submerged 2 cm below the water's surface (Fig. 1A1). Acoustic stimuli consisted of single 500 ms pure tones repeated 8 times at a rate of one every 1.5 s. Acoustic stimuli were randomly presented at the following frequencies 105, 125, 145, 165, 185, 205, 245, 285, 305, 405, 505, 605, 705, 805, 905, and 1,005 Hz, which encompasses the dominant bandwidth frequencies contained within the male midshipman advertisement call and avoids frequencies that could potentially cause interference associated with resonance frequencies of the experimental tank [see Rogers and Sisneros (29) for tank acoustic properties].

All acoustic stimuli were calibrated relative to the stimuli's sound pressure (dB re:  $1 \mu\text{Pa}$ ) via a mini-hydrophone (model 8103, Bruel and Kjaer, Naerum, Denmark) connected to a conditioning amplifier (gain = 100 mV/Pa, Nexis 2692-OS1, Bruel and Kjaer, Naerum, Denmark). However, only certain groups of fishes can detect sound pressure via secondary structures that are close in proximity or connect to the inner ear, and function to convert the received sound pressure wave into local particle motion that stimulates the inner ear. Previous midshipman studies showed that both the saccule and lagena are sound pressure sensitive based on their proximity to the swim bladder (35, 37). However, it remains to be determined if the utricle is sensitive to sound pressure; therefore, we also report the equivalent particle acceleration levels (dB re:  $1 \text{ ms}^{-2}$ ) that corresponded to the sound pressure levels (dB re:  $1 \mu\text{Pa}$ ) used in this study, based on our calibration procedures (detailed below).

Particle acceleration levels (dB re:  $1 \text{ ms}^{-2}$ ) were determined by suspending a neutrally buoyant waterproofed tri-axial accelerometer [Model VW3567A12; Sensitivity at 100 Hz:  $10.42 \text{ mV/ms}^2$  (x-axis),  $10.03 \text{ mV/ms}^2$  (y-axis),  $10.37 \text{ mV/ms}^2$  (z-axis); PCB Piezotronics, Depew, NY] that connected to a signal conditioner (gain =  $\times 100/\text{axis}$ ; Model: 482A16; PCB Piezotronics, Depew, NY). For both sound pressure (dB re:  $1 \mu\text{Pa}$ ) and particle acceleration (dB re:  $1 \text{ ms}^{-2}$ ) measurements,



**Figure 1.** Experimental design. **A:** schematic of experimental physiology tank. For all physiology experiments, fish were suspended 4 cm below the water's surface and positioned such that the face of the underwater speaker was 10 cm rostral of the otic capsule. Labels are as follows: 1) underwater speaker, 2) hydrophobic water dam, 3) parafilm suspension, 4) head holder, 5) respiration tube, 6) glass microelectrode, and 7) rocky sediment. Physiology tank dimensions: 40 cm diameter and 20 cm water depth. **B:** dorsal view of midshipman cranial cavity. The dashed circles indicate the position of the left (UL) and right (UR) inner ear utricle. **C:** cerebellum; **M:** midbrain; **SL** and **SR**, left and right saccule, respectively; **T:** telencephalon. **C:** for all experiments, pure tone acoustic stimuli (right; 500 ms duration; 8 repetitions) were delivered via an underwater speaker and evoked hair cell responses were recorded to each acoustic stimulus presentation. Consecutive ( $n = 8$ ) utricular hair cell-evoked responses (middle) were averaged (mean) across stimulus frequency, and frequency-dependent averaged output signals (right) were used to construct iso-intensity level response curves (**D**).

the mini-hydrophone and particle accelerometer, respectively, were suspended 10 cm perpendicular to the face of the underwater speaker and 4 cm below the water's surface to coincide with the position of the midshipman inner ear during auditory-evoked hair cell potential measurements. Sound pressure level (dB re: 1  $\mu$ Pa) measurements were calibrated by measuring the peak-to-peak (pk-pk) voltage ( $V_{pk-pk}$ ) amplitude on an oscilloscope (Tektronix, Beaverton, OR) and then equalized in sound pressure level (dB re: 1  $\mu$ Pa) using a custom MATLAB (MathWorks Inc., Natick, MA) script, which measured the power spectral density for all tested frequencies. The signal ( $V_{pk-pk}$ ) sent to the speaker was scaled until a reference peak-to-peak sound pressure level (SPL<sub>pk-pk</sub>) output from the speaker of  $130 \pm 0.5$  dB re: 1  $\mu$ Pa was achieved. Particle acceleration level (dB re: 1  $\text{ms}^{-2}$ ) measurements were acquired by measuring the particle motion amplitude ( $V_{pk-pk}$ ) of each tested frequency across the entire range of sound levels using a National Instruments data acquisition system (Model: NI USB-6009, National Instruments, Austin, TX) and visualized using LabVIEW software (National Instruments, Austin, TX). Using a custom LabVIEW (National Instruments, Austin, TX) script, particle motion amplitude measurements ( $V_{pk-pk}$ ) for each axis ( $x$ -,  $y$ -, and  $z$ -axis) were corrected for the gain (sensitivity) of the accelerometer. Particle motion values (dB re: 1  $\text{ms}^{-2}$ ) for each test frequency at three representative sound levels (130, 142, and 154 dB re: 1  $\mu$ Pa) are displayed in Supplemental Fig. S2.

### Utricular Potential Measurements

The methodology for recording utricular hair cell potentials follows the techniques used in our previous study, which measured the auditory-evoked potentials from the utricular hair cells of adult male plainfin midshipman (29). Midshipman were anesthetized by immersion in a 0.025% ethyl *p*-aminobenzoate (benzocaine)-buffered saltwater bath and then given an intramuscular injection of bupivacaine HCL ( $\sim 1$  mg/kg of BM) and cisatracurium besylate ( $\sim 3$  mg/kg of BM) for analgesia and immobilization, respectively. A craniotomy was then performed lateral to the sagittal crest of the skull to expose the inner ear saccule and utricle and the brain (Fig. 1B), and a hydrophobic barrier ( $\sim 2.5$  cm dia.  $\times$  5 cm height) made of denture adhesive cream (Fixodent, Proctor and Gamble Company, Cincinnati, OH) was constructed around the craniotomy to prevent saltwater contamination during experimental testing (Fig. 1A2). Fish were then transferred to the experimental tank (40 cm diameter, 20 cm water depth), which was maintained on a vibration-isolation table (TMC Vibration Control, Peabody, MA) inside a sound attenuation chamber (Industrial Acoustics, New York, NY), suspended in the center of the experimental tank using acoustically transparent film (Fig. 1A3), head-fixed 4 cm below the water's surface via a custom-built acrylic head holder (Fig. 1A4), and perfused with chilled saltwater (13–15°C) throughout experimental testing (Fig. 1A5).



Auditory-evoked utricular hair cell potentials were recorded using borosilicate glass microelectrodes (2 mm outer diameter; 1.16 mm inner diameter; A-M Systems, Sequim, WA) that were pulled using a Narishige puller (Model: PE-21) and filled with 3 M KCl (impedance: 4.0–8.0 MΩ). Electrodes were positioned in close proximity ( $\leq 2$  mm) to the medial region of the utricle near the sensory epithelia (Fig. 1A6). The analog-evoked potential signals were preamplified ( $10\times$ ; Model 5 A, Getting Instruments, San Diego, CA), bandpass filtered (0.07 to 3 kHz), and then amplified ( $10\times$ ) again via a digital filter (model SR650, Stanford Research Systems, Sunnyvale, CA). Using a lock-in amplifier (SR830, Stanford Research Systems, Sunnyvale, CA), the output signal, which was proportional to the utricular hair cell-evoked response to the stimulus fundamental frequency, was locked to a reference frequency set to the second harmonic of the pure tone stimulus frequency (i.e.,  $2 \times$  fundamental frequency), which due to populations of oppositely oriented hair cells in the teleost inner ear corresponds to the greatest evoked potential amplitudes (31, 38–40) (Fig. 1C). At the start of each experimental recording session, control trials (i.e., no sound stimulus) were conducted to measure background utricular potential levels ( $n = 8$  measurements) under ambient sound levels ( $-71 \pm 1$  dB re:  $1 \text{ ms}^{-2}$ ;  $76 \pm 1$  dB re:  $1 \mu\text{Pa}$ ). After determining background levels, stimulus trials across the experimental frequency bandwidth were carried out to construct iso-intensity level responses at various sound levels (Fig. 1D). All experimental trials were carried out using a custom MATLAB script, which controlled stimulus timing and acquired data, and all data were stored on a desktop computer.

## Analyses

Utricular hair cell auditory threshold tuning curves relative to particle acceleration (dB re:  $1 \text{ ms}^{-2}$ ) and sound pressure (dB re:  $1 \mu\text{Pa}$ ) were determined via input-output measurements of the evoked receptor potentials over the range of tested frequencies (105–1,005 Hz) and sound levels ( $-46.1$  to  $1.8$  dB re:  $1 \text{ ms}^{-2}$ ;  $103$ – $154$  dB re:  $1 \mu\text{Pa}$ ). The auditory threshold level was defined as the lowest stimulus level that yielded the lowest mean utricular-evoked potential that was at least two standard deviations above the background electrical noise measurement. The frequency that evoked the lowest utricular threshold was defined as the characteristic frequency (CF), whereas the frequency that elicited the highest evoked utricular hair cell potential response for a given sound was defined as the best frequency (BF). Particle acceleration level (dB re:  $1 \text{ ms}^{-2}$ ) thresholds were calculated as the combined magnitude vector of particle acceleration in dB scale (Eq. 1) (33, 35, 41–44) as follows:

$$\text{dB re: } 1 \text{ ms}^{-2} = 20 \text{ Log}_{10}(\sqrt{x^2 + y^2 + z^2}) \quad (1)$$


For all statistical tests, the significance level was defined at 0.05. To determine if reproductive state plays a role in modulating utricular hair cell auditory thresholds, the effects of reproductive state and stimulus frequency were analyzed via a repeated-measures analysis of variance (ANOVA, between-subject factor: reproductive state, within-subject factor: frequency  $\times$  reproductive state). As we were only interested in

how reproductive state modulates frequency sensitivity, a priori pairwise  $t$  tests compared the frequency-dependent auditory sensitivity of females from different reproductive states at the same frequency across the stimulus frequency bandwidth (105–1,005 Hz). In addition, separate two-sample  $t$  tests were performed to determine significant differences between the SL, BM, and GSI of reproductive and nonreproductive fish. All statistical analyses were performed using MATLAB software (MathWorks Inc., Natick, MA).

## RESULTS

Auditory-evoked potentials were recorded from the utricle of 33 adult female plainfin midshipman fish: 16 nonreproductive females with standard lengths (SL) that ranged from 12.4 to 19.2 cm ( $15.0 \pm 2.2$  cm; means  $\pm$  SD), body masses (BM) that ranged from 27.3 to 55.9 g ( $36.1 \pm 9.0$  g), and gonadosomatic indices (GSI) that ranged from 0.4 to 4.0 ( $1.8 \pm 1.1$ ), and 17 reproductive females with SL that ranged from 11.6 to 20.2 cm ( $16.2 \pm 2.4$  cm), BM that ranged from 35.5 to 111.0 g ( $78.3 \pm 16.4$  g), and GSI that ranged from 15.2 to 40.6 ( $31.8 \pm 5.9$ ). When comparing the morphometrics of nonreproductive and reproductive female plainfin midshipman, there was no difference in SL (two-sample  $t$  test,  $t_{1,31} = -1.499$ ,  $P = 0.144$ ); however, both BM (two-sample  $t$  test,  $t_{1,31} = -9.069$ ,  $P < 0.001$ ), and GSI (two-sample  $t$  test,  $t_{1,31} = -19.916$ ,  $P < 0.001$ ) were larger in the reproductive females, which is reflective of their reproductive status [i.e., gravid (full of eggs) vs. nongravid females].


Auditory-evoked potentials were recorded from utricular hair cells in response to particle acceleration and sound pressure levels that ranged from  $-46.1$  to  $1.8$  dB re:  $1 \text{ ms}^{-2}$  and  $103$  to  $154$  dB re:  $1 \mu\text{Pa}$ , respectively. Iso-level response profiles of the utricular-evoked potentials were generated from the presentation of single-tone stimuli that ranged from 105 Hz to 1,005 Hz for three sound levels:  $154$  dB re:  $1 \mu\text{Pa}$  ( $-0.8$  to  $10.8$  dB re:  $1 \text{ ms}^{-2}$ ),  $142$  dB re:  $1 \mu\text{Pa}$  ( $-12.9$  to  $-0.9$  dB re:  $1 \text{ ms}^{-2}$ ), and  $130$  dB re:  $1 \mu\text{Pa}$  ( $-25.9$  to  $-12.5$  dB re:  $1 \text{ ms}^{-2}$ ), which corresponded to a broad range of sound levels that have been recorded in the nest of calling type I males (i.e.,  $154$ – $161$  dB re:  $1 \mu\text{Pa}$ ) (29, 34) and recorded at or within 1 m of a calling type I males nest (i.e.,  $130$ – $142$  dB re:  $1 \mu\text{Pa}$ ) (34, 45, 46) [see Supplemental Fig. S2 for details regarding the frequency-specific particle acceleration levels (dB re:  $1 \text{ ms}^{-2}$ ) for the three sound pressure levels:  $154$ ,  $142$ , and  $130$  dB re:  $1 \mu\text{Pa}$ ]. Figure 2 displays representative utricular-evoked iso-level response curves of nonreproductive and reproductive females in response to the bandwidth of tested frequencies (105–1,005 Hz) at  $154$ ,  $142$ , and  $130$  dB re:  $1 \mu\text{Pa}$ . Iso-level response curves consisted of profiles that had best frequencies (BFs, defined as the frequency that evoked the greatest utricular-evoked potential magnitude at a given iso-level) ranging from 105 to 205 Hz in nonreproductive and reproductive females. Both nonreproductive and reproductive females had median BFs of 145 Hz at each of the sound levels tested, with no difference in the median BFs observed between nonreproductive and reproductive females at  $154$  dB re:  $1 \mu\text{Pa}$  ( $0.4$  dB re:  $1 \text{ ms}^{-2}$ ) (Friedman test,  $\chi^2 = 0$ ,  $\text{df} = 1$ ,  $P = 1$ ),  $142$  dB re:  $1 \mu\text{Pa}$  ( $-11.6$  dB re:  $1 \text{ ms}^{-2}$ ) (Friedman test,  $\chi^2 = 0.2$ ,  $\text{df} = 1$ ,  $P = 0.6547$ ), and  $130$  dB re:  $1 \mu\text{Pa}$  ( $-23.6$  dB re:  $1 \text{ ms}^{-2}$ ) (Friedman test,  $\chi^2 = 0$ ,  $\text{df} = 1$ ,  $P = 1$ ).



**Figure 2.** Representative examples of iso-intensity level curves recorded from utricular hair cells of nonreproductive (A) and reproductive (B) female plainfin midshipman. Iso-intensity responses were recorded in response to single tone playbacks ( $n = 8$ ) at sound pressure levels of 154 (black), 142 (gray), and 130 (white) dB re: 1  $\mu$ Pa. Data are represented as means  $\pm$  1 SD; note that some error bars are minimal, and the symbols may obscure the bars.

The magnitude of the auditory-evoked potentials recorded from utricular hair cells in response to pure tone stimuli was greater in reproductive females than in nonreproductive females. Figure 3 illustrates the mean iso-level response profiles of the evoked utricular potentials from nonreproductive and reproductive females in response to pure tones (105–1,005 Hz) at 154, 142, and 130 dB re: 1  $\mu$ Pa. Reproductive females had significantly higher evoked utricular potentials than nonreproductive females within sound levels encompassing a range of biologically relevant sound levels [one-

way repeated-measures ANOVA, between-subject factor: reproductive state at 154 dB re: 1  $\mu$ Pa ( $F_{1,912} = 235.4$ ,  $P < 0.001$ ), 142 dB re: 1  $\mu$ Pa ( $F_{1,912} = 247.0$ ,  $P < 0.001$ ), and 130 dB re: 1  $\mu$ Pa ( $F_{1,912} = 166.5$ ,  $P < 0.001$ )] and exhibited a significant interaction of frequency and reproductive state at 154 dB re: 1  $\mu$ Pa ( $F_{1,15} = 12.0$ ,  $P < 0.001$ ), 142 dB re: 1  $\mu$ Pa ( $F_{1,15} = 16.7$ ,  $P < 0.001$ ), and 130 dB re: 1  $\mu$ Pa ( $F_{1,15} = 19.5$ ,  $P < 0.001$ ). In addition, frequency-specific differences in the evoked magnitude response of the utricular hair cells were also observed between nonreproductive and reproductive



**Figure 3.** Iso-level response curves recorded from utricular hair cells of nonreproductive (black) and reproductive (gray) female plainfin midshipman in response to single tone playbacks at sound pressure levels of 154 (A), 142 (B), and 130 (C) dB re: 1  $\mu$ Pa. Data are represented as means  $\pm$  1 SE; note that some error bars are minimal, and the symbols may obscure the bars. The number of animals and records for each group is indicated in parentheses.

females within each sound level tested (a priori *t* tests for pair comparisons were used to determine frequency-specific differences in utricular potentials). The magnitudes of evoked utricular potentials were greater in reproductive females than in nonreproductive females at frequencies  $\leq 505$  Hz at 154 dB re: 1  $\mu\text{Pa}$  ( $P < 0.05$ ; see Supplemental Table S1),  $\leq 805$  Hz at 142 dB re: 1  $\mu\text{Pa}$  ( $P < 0.05$ ; see Supplemental Table S2), and  $\leq 305$  Hz at 130 dB re: 1  $\mu\text{Pa}$  ( $P < 0.05$ ; see Supplemental Table S3). The greatest evoked utricular potential magnitude change with respect to differences in reproductive state occurred at 105 Hz and 125 Hz at a sound pressure level of 130 dB re: 1  $\mu\text{Pa}$  (particle acceleration level at 105 Hz =  $-20.9$  dB re: 1  $\text{ms}^{-2}$  and 125 Hz =  $-23.2$  dB re: 1  $\text{ms}^{-2}$ ); at this sound pressure level, reproductive females had evoked potentials that were 6.3 and 6.2 times greater than in nonreproductive females, respectively (see Supplemental Table S3). In sum, reproductive females exhibited greater evoked utricular potentials than nonreproductive females across the frequency bandwidth tested here, with mean magnitudes that were 2.2, 2.7, and 4.1 times greater at sound pressure levels of 154 dB re: 1  $\mu\text{Pa}$  (for frequencies  $\leq 505$  Hz), 142 dB re: 1  $\mu\text{Pa}$  (frequencies  $\leq 805$  Hz) and 130 dB re: 1  $\mu\text{Pa}$  (frequencies  $\leq 305$  Hz), respectively (see Supplemental Tables S1, S2, and S3).

Auditory threshold curves based on particle acceleration (dB re: 1  $\text{ms}^{-2}$ ) and sound pressure (dB re: 1  $\mu\text{Pa}$ ) were constructed from the evoked utricular potential recordings. Figure 4 illustrates representative nonreproductive and reproductive female auditory threshold curves based on particle acceleration (dB re: 1  $\text{ms}^{-2}$ ) and sound pressure (dB re: 1  $\mu\text{Pa}$ ). In general, the utricular auditory threshold tuning curves of both nonreproductive and reproductive females exhibited the lowest thresholds at frequencies  $\leq 205$  Hz and steadily increased to the highest thresholds at frequencies  $\geq 705$  Hz. Characteristic frequencies (CFs, defined as the frequency that evoked the lowest utricular threshold) for nonreproductive females ranged from 105 to 205 Hz (median CF = 105 Hz and 145 Hz based on particle acceleration and sound pressure level tuning profiles, respectively), whereas for reproductive females, CFs ranged from 105 to 185 Hz (median CF = 105 Hz based on both particle acceleration and sound pressure level tuning profiles). The CFs based on particle acceleration did not differ with respect to reproductive state (Friedman test,  $\chi^2 = 0.2$ ,  $df = 1$ ,  $P = 0.6547$ ); however, the CFs based on sound pressure were lower in reproductive females than in nonreproductive females (Friedman test,  $\chi^2 = 6$ ,  $df = 1$ ,  $P = 0.01431$ ).

The threshold tuning curves of nonreproductive and reproductive females relative to particle acceleration (dB re: 1  $\text{ms}^{-2}$ ) and sound pressure (dB re: 1  $\mu\text{Pa}$ ) levels are summarized in Fig. 5. In general, for females of both reproductive states, the lowest utricular thresholds occurred at the lowest frequency tested (i.e., 105 Hz) (nonreproductive females: mean particle acceleration level threshold =  $-28.9 \pm 1.7$  dB re: 1  $\text{ms}^{-2}$ , mean sound pressure level threshold =  $121.5 \pm 1.7$  dB re: 1  $\mu\text{Pa}$ ; reproductive females: mean particle acceleration level threshold =  $-36.5 \pm 1.9$  dB re: 1  $\text{ms}^{-2}$ ; mean sound pressure level threshold =  $113.4 \pm 1.9$  dB re: 1  $\mu\text{Pa}$ ), whereas the highest auditory threshold levels occurred between 705 Hz and 1,005 Hz (nonreproductive females: mean particle acceleration level threshold range =  $-4$  to  $-1$  dB re: 1  $\text{ms}^{-2}$ , mean

sound pressure level threshold range = 150–153 dB re: 1  $\mu\text{Pa}$ ; reproductive females: mean particle acceleration level threshold range =  $-8$  to  $-3$  dB re: 1  $\text{ms}^{-2}$ ; mean sound pressure level threshold range = 148–151 dB re: 1  $\mu\text{Pa}$ ). The auditory thresholds were lower (i.e., more sensitive) in reproductive females than in nonreproductive females (one-way repeated-measures ANOVA, between-subject factor: reproductive state, particle acceleration level:  $F_{1,893} = 472.6$ ,  $P < 0.001$ , sound pressure level:  $F_{1,893} = 473.6$ ,  $P < 0.001$ ) and a significant interaction was observed between reproductive state and frequency (one-way repeated-measures ANOVA, within-subject factor: frequency \* reproductive state, particle acceleration level:  $F_{1,15} = 3.5$ ,  $P < 0.001$ , sound pressure level:  $F_{1,15} = 3.7$ ,  $P < 0.001$ ). Furthermore, frequency-specific differences in auditory thresholds were observed between nonreproductive and reproductive females with reproductive females being more sensitive than nonreproductive females at frequencies from 105 to 805 Hz (a priori *t* tests for pairwise comparisons of nonreproductive and reproductive females across frequency,  $P < 0.001$ ).


## DISCUSSION

The goal of this study was to determine whether seasonal changes in reproductive state modulate the auditory sensitivity of the utricle in female plainfin midshipman. We show that the utricular hair cells of reproductive females exhibit up to a sixfold magnitude increase in their evoked response to auditory stimuli and have particle acceleration thresholds that are 7–10 dB re: 1  $\text{ms}^{-2}$  lower (i.e., more sensitive) than nonreproductive females across a frequency bandwidth that includes the dominant frequencies contained within type I male vocalizations. To our knowledge, this is the first study to demonstrate reproductive state-dependent plasticity of the frequency sensitivity and auditory gain in the teleost utricle, an inner ear end organ not often associated with auditory function. In this discussion, we consider how changes in midshipman utricular auditory sensitivity may facilitate acoustic communication during social and reproductive behaviors.

### Auditory Sensitivity of the Midshipman Utricle

In mammals, the utricle primarily serves a vestibular function as it detects linear acceleration, senses horizontal translational movements, and plays an important role in static balance. However, in teleost fishes, the utricle is one of three inner ear otolithic end organs (along with the saccule and lagena) that acts as an inertial accelerometer and responds to direct displacement by acoustic particle motion and linear accelerations primarily in the horizontal plane (21, 47, 48). Although the saccule and lagena are most often implicated in sound detection and directional hearing (49–51), the utricle is posited to serve primarily a vestibular role functioning to detect head/body position relative to gravity (i.e., acts as a gravistatic organ) (25–27).

In our current study, we show that the female midshipman utricle, especially in the reproductive state, is sensitive to a broad range of acoustic frequencies with a relatively high gain in particle acceleration sensitivity (dB re: 1  $\text{ms}^{-2}$ ) from 105 to 1,005 Hz (Fig. 5). Our results confirm previous studies, which showed that the utricle in batrachoid fishes




**Figure 4.** Representative (*top*) particle acceleration (dB re:  $1 \text{ ms}^{-2}$ ) and (*bottom*) sound pressure (dB re:  $1 \text{ } \mu\text{Pa}$ ) level auditory threshold tuning curves recorded from nonreproductive (A; black) and reproductive (B; gray) female plainfin midshipman. Tuning curves were constructed using utricular hair cell-evoked responses, with thresholds defined as the lowest sound pressure level (dB re:  $1 \text{ } \mu\text{Pa}$ ) needed to evoke a utricular potential at least 2 SD above the background electrical noise level.

(toadfishes and midshipman) serves an auditory function and is capable of detecting behaviorally relevant acoustic stimuli (28, 29). Further support for the utricle of batrachoids serving an auditory function is the neuroanatomical evidence provided by Highstein et al. (52) and Sisneros et al. (53). Highstein et al. (52) showed that utricular afferents in

toadfish project to the rostral “finger” and dorsolateral aspect of the hindbrain descending octaval nucleus (DON), whereas Sisneros et al. (53) showed that the midshipman utricle has extensive projections to the intermediate and rostral intermediate auditory zones of the hindbrain DON; note that the rostral “fingerlike” extension described by Highstein et al.





**Figure 5.** Particle acceleration (A) (dB re:  $1 \text{ ms}^{-2}$ ) and sound pressure (B) (dB re:  $1 \text{ } \mu\text{Pa}$ ) level auditory threshold tuning curves recorded from nonreproductive (black) and reproductive (gray) female midshipman utricular hair cells. The auditory thresholds were defined as the lowest auditory stimulus level needed to evoke utricular potentials at least 2 SD above the background electrical noise level. All data are plotted as means  $\pm$  95% confidence interval. The number of animals and records for each group is indicated in parentheses.

(52) is similar in position and extent to the rostral intermediate zone of the midshipman DON, as described by Bass et al. (54). Furthermore, Sisneros et al. (53) showed via transneuronal labeling that the principal cells in the midshipman DON that receive input from utricular afferents subsequently project centrally to terminals in the auditory region of the mid-brain torus semicircularis similar to the saccule. Taken together, these physiological and neuroanatomical studies in batrachoid fishes strongly suggest that the utricle serves an auditory function and can detect biologically relevant acoustic stimuli.


### Seasonal Auditory Plasticity of the Utricle

We show that female utricular hair cells exhibit seasonal, reproductive state-dependent changes in evoked responses to auditory stimuli (Fig. 3), such that reproductive females exhibit greater evoked utricular potentials than nonreproductive females (Supplemental Tables S1, S2, and S3). The greatest difference in evoked potential magnitude relative to reproductive state occurred at 105 Hz and a sound pressure level of 130 dB re:  $1 \text{ } \mu\text{Pa}$  ( $-20.9 \text{ dB re: } 1 \text{ ms}^{-2}$ ) such that reproductive females displayed average utricular potentials that were  $\sim 6.3$  times greater than in nonreproductive females. Reproductive state-dependent changes in saccular-evoked potential magnitude have previously been examined in reproductive females, which have average evoked potentials  $\sim 7.4$  times greater than in nonreproductive females at 105 Hz and a sound pressure level of 130 dB re:  $1 \text{ } \mu\text{Pa}$  (13). One hypothesis for these changes in the magnitude of the hair cell-evoked potentials may, in part, be related to seasonal increases in hair cell density. Coffin et al. (36) showed that reproductive female midshipman exhibit a 13% increase in saccular hair cell density, which was paralleled by a dramatic increase in the magnitude of evoked saccular potentials. However, reproductive females and type I males do not exhibit reproductive state-dependent changes in the hair cell density of the utricle (36, 40), yet, reproductive females exhibit seasonal changes in the magnitude of evoked utricular potentials. Indeed, seasonal changes in saccular potential magnitude in reproductive females may still be related to the saccular-specific hair cell addition and may explain, in part, some of the evoked potential differences between the saccule and utricle (i.e., the utricle having  $\sim 6.3$ -fold increase

versus the saccule having a  $\sim 7.4$ -fold increase). An alternative, but not mutually exclusive, hypothesis for the change in the magnitude of hair cell potentials may be due to reproductive state-dependent changes in ion channel expression and the current kinetics of hair cells in the utricle and saccule (see Ref. 55). Future studies that characterize the ion channel current kinetics of hair cells in nonreproductive and reproductive females may provide insight into the mechanism responsible for the reproductive state-dependent changes in the magnitude of evoked potentials in the midshipman utricle and saccule.

Concurrent with the dramatic increase in utricular potential magnitude, we also observed a remarkable increase in the utricular auditory sensitivity of reproductive females when compared with nonreproductive females. The greatest change in utricular auditory sensitivity occurred from 105 to 505 Hz (Fig. 5), with reproductive females exhibiting particle acceleration thresholds that were 7–10 dB (re:  $1 \text{ ms}^{-2}$ ) lower than nonreproductive females (an increase in sensitivity equal to  $\sim 2$ –3 times) (Fig. 6A). This reproductive state-dependent increase in female auditory sensitivity corresponds with the dominant frequencies contained within type I male vocalizations, which include grunts, growls, and advertisement calls or “hums” (Fig. 6B). Grunts are short-duration (50–200 ms) broadband acoustic signals that are produced either individually or in a series of “trains” (Fig. 6B, bottom), whereas growls are longer-duration ( $> 1 \text{ s}$ ) broadband signals (Fig. 6B, middle). In general, these vocalizations are produced in an agonistic context to fend off potential rivals/intruders during nest defense (56). In contrast, hums are long-duration (up to 2 h in captive conditions) multiharmonic acoustic courtship signals that have fundamental frequencies ranging from 80 to 102 Hz (15, 45, 46). Compared with broadband grunts and growls, which have much of their spectral energy at frequencies  $< 600 \text{ Hz}$ , hums have prominent harmonics ranging up to  $\sim 500 \text{ Hz}$ , with additional lower amplitude harmonics ranging up to 1,000 Hz (see Fig. 6B, top). Together, our results suggest the utricle of reproductive females is better adapted than in nonreproductive females to detect the dominant spectral energy contained within midshipman social acoustic signals (hums, growls, and grunts), which correspond to frequencies  $< 600 \text{ Hz}$  (Fig. 6B). Thus, reproductive state-dependent changes in utricular





**Figure 6.** Comparison of female midshipman particle motion (dB re: 1 ms<sup>-2</sup>) sensitivity and male midshipman vocalizations. **A:** utricular hair cell particle acceleration threshold difference ( $\Delta$  dB re: 1 ms<sup>-2</sup>) between reproductive and nonreproductive female midshipman. **B:** power spectral density (dB re: 1  $\mu$ Pa<sup>2</sup>/Hz) curves of a male midshipman hum (top), growl (middle), and grunt (bottom). Inset: waveform of male midshipman hum, growl, and grunt, respectively. Scale bars represent 5, 1,000, and 50 ms, respectively. All vocalizations were recorded from a reproductive type I male midshipman housed in a large, indoor concrete tank (3 m diameter; 14.1°C) at the University of Washington Friday Harbor Laboratories. Source level recordings were made using a minihydrophone placed directly in front of the entrance of an artificial nest.

auditory sensitivity may represent an adaptive auditory plasticity that complements the saccular auditory sensitivity of reproductive female midshipman (Supplemental Fig. S3) and help facilitate midshipman social and reproductive acoustic communication.

### Potential Mechanisms for Utricular Auditory Plasticity

The observed changes in utricular auditory sensitivity are likely due to seasonal changes in circulating gonadal steroids (androgens and estrogens), which are related to seasonal

changes in midshipman reproductive state (57). Saccular afferents in nonreproductive females treated with either testosterone or 17 $\beta$ -estradiol exhibit enhanced frequency sensitivity and phase-locking accuracy to higher frequencies within the midshipman hearing range, which effectively enhances acoustic communication (20). Concurrent with reproductive state-dependent changes in gonadal steroid levels are parallel changes in the large-conductance, calcium-activated potassium (BK) channels, which are responsible for the rapid outward currents that contribute to the electrical resonance and low-frequency (<1 kHz) tuning of hair cells in nonmammalian vertebrates (58–60). Rohmann et al. (55) demonstrated that saccular hair cells of reproductive midshipman exhibit increased expression of calcium-activated BK channels, which is correlated with enhanced higher frequency sensitivity (>145 Hz) and that pharmacological inhibition of BK channels results in decreased saccular sensitivity similar to nonreproductive fish. Together, these studies suggest that gonadal steroids may modulate seasonal changes in frequency sensitivity via the regulation of hair cell BK channel expression to effectively enhance auditory sensitivity for social acoustic communication.

In addition, reproductive state-dependent changes in dopaminergic efferent projections to the inner ear may also be responsible for the observed seasonal, reproductive state-dependent changes in utricular sensitivity. Previous work by Forlano et al. (17) showed that dopaminergic innervation of the saccule varied with reproductive state such that reproductive females have a seasonal reduction in dopaminergic input. Furthermore, Perelmuter et al. (61) showed that dopamine decreases saccular auditory sensitivity via a D2-like receptor and that D2a receptor expression is reduced in the saccule during the midshipman breeding season. Perelmuter et al. (61) also found that saccular auditory sensitivity is modulated by the dopaminergic efferent system, whereby a release in inhibition effectively mimics the reproductive auditory phenotype and enhances peripheral encoding of social acoustic signals. Furthermore, Perelmuter et al. (62) recently showed that testosterone treatment mimics the seasonal downregulation of dopamine in the midshipman saccule, which provides evidence that steroid regulation of peripheral auditory sensitivity is mediated, at least in part, by dopamine. Future studies that examine similar reproductive state-dependent, gonadal steroid regulatory mechanisms of hair cell ion channel expression and dopaminergic innervation to the utricle will be instrumental in understanding the neuroendocrine basis of peripheral auditory sensitivity modulation in midshipman fish and other vertebrates, including mammals.

### Conclusions

The utricle in mammals primarily serves as a vestibular organ for detecting linear acceleration and sensing translational movements in the horizontal plane. However, in fishes, the utricle is one of three inner ear otolithic end organs (saccule, utricle, and lagena) that act as biological accelerometers and respond to acoustic particle motion and horizontal linear accelerations. Although, to some degree, all three otolithic end organs in teleost fishes are posited to

serve both an auditory and vestibular function, the teleost utricle is often thought to primarily serve a vestibular function. Here, we show that the utricle in the vocal plainfin midshipman serves an auditory function that is seasonally plastic and modulated by the animal's reproductive state, effectively enhancing the utricle's auditory sensitivity to conspecific acoustic signals. Whether these seasonal-dependent changes extend beyond the auditory system to the vestibular system has yet to be assessed and should be considered in future vestibular research, given the multimodal function of the inner ear end organs.

## SUPPLEMENTAL DATA

Supplemental Figs. S1–S3 and Supplemental Tables S1–S3: <https://doi.org/10.6084/m9.figshare.20363625>.

## ACKNOWLEDGMENTS

The authors thank the captain and crew of the R/V Kittiwake for assistance with winter fish collection, and Sujay Balebail and Lorin Gardner for assistance with summer fish collection and fish husbandry.

## GRANTS

This work was supported by a National Science Foundation grant to J.A.S. (IOS 1933166) and A.B.C. (IOS 1932898). L.S.R. was supported by a National Science Foundation Graduate Research Fellowship (DGE 1762114), National Institutes of Health auditory neuroscience training grant (T32DC005361), and a Howard Hughes Medical Institute Gilliam Fellowship (GT15044).

## DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

## AUTHOR CONTRIBUTIONS

L.S.R., A.B.C., and J.A.S. conceived and designed research; L.S.R. performed experiments; L.S.R. analyzed data; L.S.R. and J.A.S. interpreted results of experiments; L.S.R. prepared figures; L.S.R. and J.A.S. drafted manuscript; L.S.R., A.B.C., and J.A.S. edited and revised manuscript; L.S.R., A.B.C., and J.A.S. approved final version of manuscript.

## REFERENCES

- Ball GF, Castelino CB, Maney DL, Appeltants D, Balthazart J. The activation of birdsong by testosterone: multiple sites of action and role of ascending catecholamine projections. *Ann N Y Acad Sci* 1007: 211–231, 2003. doi:10.1196/annals.1286.021.
- Bass AH, Zakon HH. Sonic and electric fish: at the crossroads of neuroethology and behavioral neuroendocrinology. *Horm Behav* 48: 360–372, 2005. doi:10.1016/j.yhbeh.2005.05.022.
- Wilczynski W, Lynch KS, O'Bryant EL. Current research in amphibians: studies integrating endocrinology, behavior, and neurobiology. *Horm Behav* 48: 440–450, 2005. doi:10.1016/j.yhbeh.2005.06.001.
- Caras ML, Brenowitz E, Rubel EW. Peripheral auditory processing changes seasonally in Gambel's white-crowned sparrow. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 196: 581–599, 2010. doi:10.1007/s00359-010-0545-1.
- Caras ML, O'Brien M, Brenowitz EA, Rubel EW. Estradiol selectively enhances auditory function in avian forebrain neurons. *J Neurosci* 32: 17597–17611, 2012. doi:10.1523/JNEUROSCI.3938-12.2012.
- Caras ML, Sen K, Rubel EW, Brenowitz EA. Seasonal plasticity of precise spike timing in the avian auditory system. *J Neurosci* 35: 3431–3445, 2015. doi:10.1523/JNEUROSCI.3407-14.2015.
- Gall MD, Salameh TS, Lucas JR. Songbird frequency selectivity and temporal resolution vary with sex and season. *Proc Biol Sci* 280: 20122296, 2013. doi:10.1098/rspb.2012.2296.
- Henry KS, Lucas JR. Vocationally correlated seasonal auditory variation in the house sparrow (*Passer domesticus*). *J Exp Biol* 212: 3817–3822, 2009. doi:10.1242/jeb.033035.
- Vélez A, Gall MD, Lucas JR. Seasonal plasticity in auditory processing of the envelope and temporal fine structure of sounds in three songbirds. *Animal Behaviour* 103: 53–63, 2015. doi:10.1016/j.anbehav.2015.01.036.
- Goense JBM, Feng AS. Seasonal changes in frequency tuning and temporal processing in single neurons in the frog auditory midbrain. *J Neurobiol* 65: 22–36, 2005. doi:10.1002/neu.20172.
- Miranda JA, Wilczynski W. Female reproductive state influences the auditory midbrain response. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 195: 341–349, 2009. doi:10.1007/s00359-008-0410-7.
- Penna M, Capranica RR, Somers J. Hormone-induced vocal behavior and midbrain auditory sensitivity in the green treefrog, *Hyla cinerea*. *J Comp Physiol A* 170: 73–82, 1992. doi:10.1007/BF00190402.
- Sisneros JA. Seasonal plasticity of auditory saccular sensitivity in the vocal plainfin midshipman fish, *Porichthys notatus*. *J Neurophysiol* 102: 1121–1131, 2009. doi:10.1152/jn.00236.2009.
- Sisneros JA, Bass AH. Seasonal plasticity of peripheral auditory frequency sensitivity. *J Neurosci* 23: 1049–1058, 2003. doi:10.1523/jneurosci.23-03-01049.2003.
- Bass AH, Bodnar DH, Marchaterre MA. Complementary explanations for existing phenotypes in an acoustic communication system. In: *The Design of Animal Communication*, edited by Hauser MD, Konishi M. Cambridge, MA: MIT Press, 1999, p. 493–514.
- Bass AH, McKibben JR. Neural mechanisms and behaviors for acoustic communication in teleost fish. *Prog Neurobiol* 69: 1–26, 2003. doi:10.1016/S0301-0082(03)00004-2.
- Forlano PM, Ghahramani ZN, Monestime CM, Kurochkin P, Chernenko A, Milkis D. Catecholaminergic innervation of central and peripheral auditory circuitry varies with reproductive state in female midshipman fish, *Porichthys notatus*. *PLoS One* 10: e0121914, 2015. doi:10.1371/journal.pone.0121914.
- Sisneros JA. Adaptive hearing in the vocal plainfin midshipman fish: getting in tune for the breeding season and implications for acoustic communication. *Integr Zool* 4: 33–42, 2009. doi:10.1111/j.1749-4877.2008.00133.x.
- Forlano PM, Maruska KP, Sisneros JA, Bass AH. Hormone-dependent plasticity of auditory systems in fishes. In: *Hearing and Hormones*, edited by Bass AH, Sisneros JA, Popper AN, Fay RR. Cham, Switzerland: Springer International Publishing, 2016, p. 15–51.
- Sisneros JA, Forlano PM, Deitcher DL, Bass AH. Steroid-dependent auditory plasticity leads to adaptive coupling of sender and receiver. *Science* 305: 404–407, 2004. doi:10.1126/science.1097218.
- Popper AN, Fay RR. Sound detection and processing by fish: critical review and major research questions. *Brain Behav Evol* 41: 14–38, 1993. doi:10.1159/000113821.
- Popper AN, Fay RR. Rethinking sound detection by fishes. *Hear Res* 273: 25–36, 2011. doi:10.1016/j.heares.2009.12.023.
- Bianco IH, Ma L-H, Schoppik D, Robson DN, Orger MB, Beck JC, Li JM, Schier AF, Engert F, Baker R. The tangential nucleus controls a gravito-inertial vestibulo-ocular reflex. *Curr Biol* 22: 1285–1295, 2012. doi:10.1016/j.cub.2012.05.026.
- Boyle R, Mensinger AF, Yoshida K, Usui S, Intravaia A, Tricas T, Highstein SM. Neural readaptation to Earth's gravity following return from space. *J Neurophysiol* 86: 2118–2122, 2001. doi:10.1152/jn.2001.86.4.2118.
- Boyle R, Popova Y, Varelas J. Influence of magnitude and duration of altered gravity and readaptation to 1 g on the structure and function of the utricle in toadfish, *Opsanus tau*. *Front Physiol* 9: 1469, 2018. doi:10.3389/fphys.2018.01469.
- Inoue M, Tanimoto M, Oda Y. The role of ear stone size in hair cell acoustic sensory transduction. *Sci Rep* 3: 2114, 2013. doi:10.1038/srep02114.

27. **Riley BB, Moorman SJ.** Development of utricular otoliths, but not saccular otoliths, is necessary for vestibular function and survival in zebrafish. *J Neurobiol* 43: 329–337, 2000. doi:10.1002/1097-4695(20000615)43:4<3C329::aid-neu2%3E3.0.co;2-h.
28. **Maruska KP, Mensinger AF.** Directional sound sensitivity in utricular afferents in the toadfish *Opsanus tau*. *J Exp Biol* 218: 1759–1766, 2015. doi:10.1242/jeb.115345.
29. **Rogers LS, Sisneros JA.** Auditory evoked potentials of utricular hair cells in the plainfin midshipman, *Porichthys notatus*. *J Exp Biol* 223: jeb226464, 2020. doi:10.1242/jeb.226464.
30. **Rogers LS, Van Wert JC, Mensinger AF.** Response of toadfish (*Opsanus tau*) utricular afferents to multimodal inputs. *J Neurophysiol* 128: 364–377, 2022. doi:10.1152/jn.00483.2021.
31. **Sisneros JA.** Saccular potentials of the vocal plainfin midshipman fish, *Porichthys notatus*. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 193: 413–424, 2007. doi:10.1007/s00359-006-0195-5.
32. **Alderks PW, Sisneros JA.** Ontogeny of auditory saccular sensitivity in the plainfin midshipman fish, *Porichthys notatus*. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 197: 387–398, 2011. doi:10.1007/s00359-010-0623-4.
33. **Bhandiwad AA, Whitchurch EA, Colleye O, Zeddies DG, Sisneros JA.** Seasonal plasticity of auditory saccular sensitivity in “sneaker” type II male plainfin midshipman fish, *Porichthys notatus*. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 203: 211–222, 2017. doi:10.1007/s00359-017-1157-9.
34. **Vetter BJ, Seeley LH, Sisneros JA.** Lagenar potentials of the vocal plainfin midshipman fish, *Porichthys notatus*. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 205: 163–175, 2019. doi:10.1007/s00359-018-01314-0.
35. **Colleye O, Vetter BJ, Mohr RA, Seeley LH, Sisneros JA.** Sexually dimorphic swim bladder extensions enhance the auditory sensitivity of female plainfin midshipman fish, *Porichthys notatus*. *J Exp Biol* 222: jeb204552, 2019. doi:10.1242/jeb.204552.
36. **Coffin AB, Mohr RA, Sisneros JA.** Saccular-specific hair cell addition correlates with reproductive state-dependent changes in the auditory saccular sensitivity of a vocal fish. *J Neurosci* 32: 1366–1376, 2012. doi:10.1523/JNEUROSCI.4928-11.2012.
37. **Vetter BJ, Sisneros JA.** Swim bladder enhances lagenar sensitivity to sound pressure and higher frequencies in female plainfin midshipman (*Porichthys notatus*). *J Exp Biol* 223: jeb225177, 2020. doi:10.1242/jeb.225177.
38. **Cohen MJ, Winn HE.** Electrophysiological observations on hearing and sound production in the fish, *Porichthys notatus*. *J Exp Zool* 165: 355–369, 1967. doi:10.1002/jez.1401650305.
39. **Furukawa T, Ishii Y.** Neurophysiological studies on hearing in goldfish. *J Neurophysiol* 30: 1377–1403, 1967. doi:10.1152/jn.1967.30.6.1377.
40. **Lozier NR, Sisneros JA.** Reproductive-state dependent changes in saccular hair cell density of the vocal male plainfin midshipman fish. *Hear Res* 383: 107805, 2019. doi:10.1016/j.heares.2019.107805.
41. **Rogers LS, Putland RL, Mensinger AF.** The effect of biological and anthropogenic sound on the auditory sensitivity of oyster toadfish, *Opsanus tau*. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 206: 1–14, 2020. doi:10.1007/s00359-019-01381-x.
42. **Vasconcelos RO, Fonseca PJ, Amorim MCP, Ladich F.** Representation of complex vocalizations in the Lusitanian toadfish auditory system: evidence of fine temporal, frequency and amplitude discrimination. *Proc Biol Sci* 278: 826–834, 2011. doi:10.1098/rspb.2010.1376.
43. **Vetter BJ.** Role of the Lagenar in fish hearing and its susceptibility to anthropogenic noise. *Proc Mtgs Acoust* 37: 010001, 2019. doi:10.1121/2.0001031.
44. **Wysocki LE, Codarin A, Ladich F, Picciulin M.** Sound pressure and particle acceleration audiograms in three marine fish species from the Adriatic Sea. *J Acoust Soc Am* 126: 2100–2107, 2009. doi:10.1121/1.3203562.
45. **Balebail S, Sisneros JA.** Long duration advertisement calls of nesting male plainfin midshipman fish are honest indicators of size and condition. *J Exp Biol* 225: jeb243889, 2022. doi:10.1242/jeb.243889.
46. **Feng NY, Bass AH.** “Singing” fish rely on circadian rhythm and melatonin for the timing of nocturnal courtship vocalization. *Curr Biol* 26: 2681–2689, 2016. doi:10.1016/j.cub.2016.07.079.
47. **De Vries H.** The mechanics of the labyrinth otoliths. *Acta Otolaryngol* 38: 262–273, 1951. doi:10.3109/00016485009118384.
48. **Fay RR.** The goldfish ear codes the axis of acoustic particle motion in three dimensions. *Science* 225: 951–954, 1984. doi:10.1126/science.6474161.
49. **Hawkins AD, Popper AN.** Directional hearing and sound source localization by fishes. *J Acoust Soc Am* 144: 3329, 2018. doi:10.1121/1.5082306.
50. **Sand O.** Directional sensitivity of microphonic potentials from the perch ear. *J Exp Biol* 60: 881–899, 1974. doi:10.1242/jeb.60.3.881.
51. **Sisneros JA, Rogers PH.** Directional hearing and sound source localization in fishes. In: *Fish Hearing and Bioacoustics: An Anthology in Honor of Arthur N. Popper and Richard R. Fay*, edited by Sisneros JA. Cham, Switzerland: Springer International Publishing, 2016, p. 121–155.
52. **Highstein SM, Kitch R, Carey J, Baker R.** Anatomical organization of the brainstem octavolateralis area of the oyster toadfish, *Opsanus tau*. *J Comp Neurol* 319: 501–518, 1992. doi:10.1002/cne.903190404.
53. **Sisneros JA, Marchaterre MA, Bass AH.** Otolithic endorgan projections of the inner ear in a vocal fish. *Bioacoustics* 12: 137–139, 2012. doi:10.1080/09524622.2002.9753674.
54. **Bass AH, Bodnar DA, Marchaterre MA.** Midbrain acoustic circuitry in a vocalizing fish. *J Comp Neurol* 419: 505–531, 2000. doi:10.1002/(sici)1096-9861(20000417)419:4<3C505::aid-cne7%3E3.0.co;2-3.
55. **Rohmann KN, Fergus DJ, Bass AH.** Plasticity in ion channel expression underlies variation in hearing during reproductive cycles. *Curr Biol* 23: 678–683, 2013. doi:10.1016/j.cub.2013.03.014.
56. **Sisneros JA.** Steroid-dependent auditory plasticity for the enhancement of acoustic communication: recent insights from a vocal teleost fish. *Hear Res* 252: 9–14, 2009. doi:10.1016/j.heares.2008.12.007.
57. **Sisneros JA, Forlano PM, Knapp R, Bass AH.** Seasonal variation of steroid hormone levels in an intertidal-nesting fish, the vocal plainfin midshipman. *Gen Comp Endocrinol* 136: 101–116, 2004. doi:10.1016/j.ygcen.2003.12.007.
58. **Fettiplace R, Fuchs PA.** Mechanisms of hair cell tuning. *Annu Rev Physiol* 61: 809–834, 1999. doi:10.1146/annurev.physiol.61.1.809.
59. **Lewis RS, Hudspeth AJ.** Voltage- and ion-dependent conductances in solitary vertebrate hair cells. *Nature* 304: 538–541, 1983. doi:10.1038/304538a0.
60. **Roberts WM, Howard J, Hudspeth AJ.** Hair cells: transduction, tuning, and transmission in the inner ear. *Annu Rev Cell Biol* 4: 63–92, 1988. doi:10.1146/annurev.cb.04.110188.000431.
61. **Perelmuter JT, Wilson AB, Sisneros JA, Forlano PM.** Forebrain dopamine system regulates inner ear auditory sensitivity to socially relevant acoustic signals. *Curr Biol* 29: 2190–2198.e3, 2019. doi:10.1016/j.cub.2019.05.055.
62. **Perelmuter JT, Hom KN, Mohr RA, Demis L, Kim S, Chernenko A, Timothy M, Middleton MA, Sisneros JA, Forlano PM.** Testosterone treatment mimics seasonal downregulation of dopamine innervation in the auditory system of female midshipman fish. *Integr Comp Biol* 61: 269–282, 2021. doi:10.1093/icb/ibab070.